

*Cyclosperrum leptophyllum* (Pers.) Britt. & P. Wilson, MARSH PARSLEY. Annual, taprooted, short-lived rosetted, 1–several-stemmed at base, branched throughout with forks forming at most canopy nodes, decumbent to erect, 5–60 cm tall; shoots with basal and cauline leaves, glabrous, with a faint, carrotlike (*Daucus*) smell. **Stems:** cylindric, to 3 mm diameter, smooth, green-striped with light green to yellowish green veins. **Leaves:** helically alternate, 3–4-pinnately compound or dissected, petiolate, without stipules; petiole sheathing stem node, to 40 mm long (basal and lower cauline leaves) or 2–10 mm long (upper cauline leaves), channeled above, rounded on lower side, with membranous wings of larger leaves 2–10 × 3.5 mm wide at base, on cauline leaves < 2 mm long; blade of basal and lower cauline leaves ovate in outline, 30–100 × 25–35 mm, typically with opposite primary and secondary lateral leaflets, ultimate segments channeled, 2–5 × 0.5–1 mm, entire; blade of upper cauline leaves 25–55 × 40–80 mm long, 3-dimensional, with ± linear segments diverging acutely from axis, ultimate segments channeled, 3–18(–30) × 0.3–0.7 mm. **Inflorescence:** compound umbel, appearing opposite leaves, of 2–3(–5) open umbellets, 6–25-flowered, typically < 25(–35) mm long, umbellet several-flowered, lacking bracts, glabrous; **involucre** absent; primary rays subequal and divergent, 8–20 mm long; involucre absent; pedicel divergent and ascending, thin, 3–6.5(–16) mm long, the central pedicels infrequent and much shorter (central flower subsessile). **Flower:** bisexual, radial, 0.8–1 mm across; **sepals** absent; **petals** 5, spreading but with tip curved inward, oblong in outline, 0.35–0.5 mm long, white or sometimes pinkish or purplish red at tip or margin, acute at tip, lacking midridge on upper (inner) surface; **stamens** 5, free, ± 0.3 mm long; filaments spreading, tapered to tip, < 0.2 mm long, whitish; anthers dorsifixed, dithecal, 0.15 mm long, light yellow becoming purplish red, longitudinally dehiscent; pollen pale yellow; **pistil** 1; ovary inferior, 2-lobed with a 2-lobed platform (stylopodium, stylar tissue) on top, oblong compressed side-to-side, 0.5–0.7 mm long and wide, strongly ribbed, ribs 10 but those next to fruit axis shorter, stylopodium greenish, 2-chambered, each chamber with 1 ovule; styles ca. 0.1 mm long. **Fruit:** schizocarp, of 2 dry, 1-seeded halves (mericarps), mericarps semi-ovate and 5-ribbed on broad sides, elliptic to oblong in edge view, flattish or slightly concave on inner face on fruit axis, (1.3–)1.8–2(–3) × 1.5–2 mm, typically brownish purple, the ribs prominent, corky, subequal in width but lateral ones shorter in length. Late November–early July.

Naturalized. Annual appearing occasionally as a weed in cultivated areas of Bel Air, Westwood, and Santa Monica adjacent to SMM, but since 2003 collected at the Chatsworth Reservoir (SH) and occasionally found along roadsides on the coastal side of SMM. *Cyclosperrum leptophyllum* most closely resembles *Apiastrum angustifolium*, which also has tiny white flowers, but can be easily distinguished by the mericarps and smell of the foliage.

B. A. Prigge & A. C. Gibson